

2602 S. 24th Street . Phoenix, AZ 85034 Tel 602 759 1245 . Fax 602 759 8331 . www.CyrexLabs.com ACCESSION #: 17-108497
REQUISITION #: T05170963
SAMPLE TYPE: Serum
DOCTOR / PATIENT ID:

PAGES: 1 of 1

DATE COLLECTED: 5/11/2017
DATE RECEIVED: 5/11/2017
DATE OF REPORT: 5/12/2017

PAGES: 1 of 1

## PRACTITIONER

## **DEMODR, DEMODR TEST**

2602 S. 24th Street Phoenix, Arizona 85034

## **PATIENT**

Name: REPORT, SAMPLE

**DOB:** 11/01/1990

Gender: M

TEST	RESULT			
Array 5 – Multiple Autoimmune Reactivity Screen **	IN RANGE (Normal)	EQUIVOCAL*	OUT OF RANGE	REFERENCE (ELISA Index)
Parietal Cell + ATPase	0.85			0.1-1.4
Intrinsic Factor	0.50			0.1-1.2
ASCA + ANCA	0.50			0.2-1.4
Tropomyosin	0.94			0.1-1.5
Thyroglobulin	0.70			0.1-1.3
Thyroid Peroxidase	0.70			0.1-1.3
21-Hydroxylase (Adrenal Cortex)	0.76			0.2-1.2
Myocardial Peptide	0.98			0.1-1.5
Alpha-Myosin	0.96			0.3-1.5
Phospholipid	0.85			0.2-1.3
Platelet Glycoprotein	0.85			0.1-1.3
Ovary/Testis ***	0.79			0.1-1.2
Fibulin	1.01			0.4-1.6
Collagen Complex	0.77			0.2-1.6
Arthritic Peptide	0.90			0.2-1.3
Osteocyte	1.09			0.1-1.4
Cytochrome P450 (Hepatocyte)	0.75			0.3-1.6
Insulin + Islet Cell	0.80			0.4-1.7
Glutamic Acid Decarboxylase 65	0.83			0.2-1.6
Myelin Basic Protein	0.81			0.1-1.4
Asialoganglioside	0.75			0.1-1.4
Alpha-Tubulin + Beta-Tubulin	0.71			0.4-1.4
Cerebellar	1.06			0.2-1.4
Synapsin	0.84			0.1-1.2

<sup>\*\*\*</sup> Ovary and Testis are tested together to avoid any confusion arising out of potential cross-reactivity

Mark G. Kartub, M.D., Medical Director

Cyrex Laboratories is certified under the Clinical Laboratory Improvement Amendments of 1988 ("CLIA") as qualified to perform high-complexity clinical testing. Test result data on its own does not constitute a diagnosis. Only a physician or qualified healthcare professional should interpret the significance of a clinical lab test or make a diagnosis. This test was developed and its performance characteristics determined by Cyrex Laboratories, LLC. The names and titles of tests and arrays are for reference purposes only.

<sup>\*\*</sup> All analytes are tested for IgG and IgA combined.

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.